



INSECTICIDE



ADAMA

# CORMORAN®

A multi-mode of action foliar insecticide for rapid knockdown and extended residual control of all damaging life stages of pests in an easy-to-use formulated mixture.



GROUPS 4 & 15

## Active Ingredients:

Novaluron 100 g/L, and Acetamiprid 80 g/L

## Packaging:

Case: 2 jugs x 10.08 L

## Application Rates and Area Treated

Rate: 440 – 750 ml/ha

Area Treated: 13 – 23 ha/jug

## Water Volume

Ground: 200 L/ha

(50 US gal/ha)

Aerial: Do not apply by air.

## Rainfastness

N/A

## SERIOUS INSECT PROTECTION FOR VEGETABLE CROPS

### REGISTERED VEGETABLE CROPS

- Brassica leafy greens (Crop Subgroup 4-13B)
- Head and stem brassica vegetable crops (Crop group 5-13)
- Peppers (bell and non-bell)
- Sweet corn

### KEY INSECTS CONTROLLED (IN VEGETABLE CROPS)

- Alfalfa looper
- Aphids
- Armyworm
- Cabbage looper
- Colorado potato beetle
- Diamondback moth
- European corn borer
- Imported cabbageworm
- Japanese beetle
- Lygus bug
- Swede midge

NOTE: CORMORAN® is also registered for use in apples, bushberries (crop subgroup 13-07B), strawberries and stone fruit (crop group 12-09) for control of more than 25 insect pests such as cherry fruit fly, green fruitworm, mullein bug, plum curculio, spotted-wing drosophila and strawberry clipper weevil.

Please refer to the label for the complete list of registered crops, pests, and associated application rates



# CORMORAN®

## CROP STAGING AND RATES

Crop	Insects Controlled	Rate (ml/ha)*	Application Instructions
Peppers (bell and non-bell)	Colorado potato beetle (CPB)	440 – 700	Do not make applications less than 7 days apart.
	Aphids	490	
	European corn borer	650 – 750	Do not apply more than 2.63 L/ha per season.  For CPB, do not apply more than twice to a single generation and do not apply to successive generations.
	Armyworm, Cabbage looper	440 – 750	
Brassica (cole) leafy vegetables (Crop Group 5-13):  Broccoli, Chinese broccoli, Broccoli raab (rapini), Brussels sprouts, Cabbage, Chinese cabbage (bok choy or napa), Chinese mustard cabbage (gai choy), Cauliflower, Cavalo broccoli, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens and cultivars, varieties and/or hybrids of these commodities	Alfalfa looper, Armyworm, Cabbage looper, Diamondback moth, Imported cabbageworm	440 – 750	Do not apply more than 1.68 L/ha per season.  Do not apply more than once every 7 days.
	Aphids	650 – 750	
	Lygus bug, Swede midge	740	
Leafy vegetables – Brassica leafy greens (Crop Subgroup 4-13B):  Arugula, Broccoli raab, Chinese broccoli, Abyssinian cabbage, Seakale cabbage, Bok choy (Chinese cabbage), Collards, Garden cress, Upland cress, Hanover salad, Kale, Maca, Mizuna, Mustard greens, Radish leaves, Rape greens, Wild rocket, Shepherd's purse, Turnip greens, Watercress, as well as cultivars, varieties and hybrids of these commodities	Aphids	650 – 750	Applications per season: 3  Do not apply more than once every 7 – 10 days.
Sweet corn	Aphids	500–700	Applications per season: 2  Do not apply more than once every 21 days.

\* Use higher rate for larger infestations.

NOTE: For all vegetable crops, apply in a minimum finished spray volume of 200 L/ha by ground.

### PRE-HARVEST INTERVALS

Brassica (cole) leafy vegetables (Crop Group 5-13),

Leafy vegetables – Brassica leafy greens

(Crop Subgroup 4-13B), Peppers: 7 days

Sweet Corn: 10 days



Always read and follow registered product label instructions.  
1.855.264.6262 ADAMA.COM

© CORMORAN is a trademark of ADAMA Agricultural Solutions Canada Ltd. All other products mentioned are trademarks of their respective companies.  
© 2022 Adama Agricultural Solutions Canada Ltd.